

EDITORIAL

**THE EFFECT OF AREA AND SITE OF TYMPANIC MEMBRANE
PERFORATIONS ON HEARING THRESHOLD AMONG SUDANESE
PATIENTS**

Abu sufian Hassan Ahmed Elhaj; M.B.B.S , F.S.M.S.B (otolaryngology)¹, Mohammad Bushara Abdalla M.B.B.S,
MM.S.A,Ph.D², Haydar Abubakar Abdalla M.B.B.S,F.R.C.S (otolaryngology)³

1-Dr. Abu sufian Hassan Ahmed Elhaj M.B .B. S, F.S.M.S.B (otolaryngology) Assistant professor, Department of surgery; University of Gezira .

2-Dr. Mohammad Bushara Abdalla M.B.B.S, MM.S.A, PhD , senior audiologist, Khartoum ENT Teaching Hospital.

3-Dr. Haydar Abubakar Abdalla M. B.B.S ,F.R.C.S (otolaryngology) Assistant professor , Department of surgery; University of Khartoum

Correspondence:

Dr. Abu sufian Hassan Ahmed Elhaj, MD(otolaryngology). Department of surgery, University of Gezira,
E.mail: abusufianelhaj@yahoo.com.

تأثير مساحة وموقع ثقب طبلة الأذن على درجة السمع

مقدمة:- التهاب الأذن الوسطى المزمن من الأمراض الأذنية الأكثر شيوعاً وسط الأطفال والكبار في السودان من العلامات المرضية له ثقب طبلة الأذن ونجيج الأذن ودرجات متفاوتة من ضعف السمع.

أهداف البحث:- يهدف هذا البحث دراسة أثر مساحة و موقع ثقب طبلة الأذن على درجة السمع لدي مرضي سودانيين.
المرضي وطريقة البحث:- هذه دراسة مستقبلية أجريت بقسمي الأذن و الأنف و الحنجرة بكل من مستشفى الخرطوم وابن سينا التعليمي في الفترة من 1 ابريل إلى 30 يوليو 2002. جمعت المعلومات من أشخاص شاهدين والمرضي بواسطة استبانة. حوت الدراسة على (25) شخص شاهد (50) إذن تم اختيارهم عشوائياً تم قياس سمعهم من اجل إيجاد مرجع لمعدل السمع الطبيعي. تم اختيار (71) من المرضي تباعا يعانون من التهاب الأذن الوسطى المزمن ، بإذن واحده أو الإذنين(100اذن) .

EDITORIAL

النتائج: - متوسط معدل درجة سمع الأصحاء هو (17) ديسبل وهو في مدى المعدل الطبيعي العالمي لذي نورثين وداون 0 – 25 ديسبل. تم إيجاد ذلك بحساب متوسط درجة الكثافة السمعيه في الترددات (500-1000-2000) هيرز. وجد أن درجة السمع في الترددات المنخفضة تتأثر أولاً بغض النظر عن موقع ومساحة ثقب الطبل وبزيادة مساحة الثقب تتأثر الترددات المتوسطة و العالية . أكثر المرضى (79%) يعانون من ضعف السمع التوصيلي . الموقع الأكثر شيوعاً لثقب طبلة الأذن هو الأوسط (90%) يليه، الثقب المتوسط الخلفي 3% . المساحة الأكثر شيوعاً لثقب الطبلة هو تحت التام (54%) يليه الثقب الكبير (25%). وجد ان نقص درجة السمع اكثر تأثيراً في الثقب الخلفي السفلي والامامي العلوي.

الخلاصة والتوصيات:-

- من هذه الدراسة نستخلص أنه بغض النظر عن مساحة ثقب طبلة الأذن نجد أن موقع ثقب طبلة الأذن هو الأكثر تأثيراً علي درجة السمع. الثقب الخلفي السفلي والامامي العلوي لهم التأثير الاكثر علي درجة السمع. من اجل هذا السبب نوصي بان المرضي الذين لديهم هذه الثقوب يجب أن تجري لهم عملية ترميم طبلة الأذن.

ABSTRACT

Background: Chronic otitis media is one of the commonest otological problems among Sudanese children and adults. The disease presents with perforations of the eardrum, discharging ears and varying degrees of hearing loss.

Objective: The objective of this work is to study the effect of the area and the site of the tympanic membrane perforation on hearing threshold among Sudanese patients.

Patients & Methods: This is a prospective hospital-based study conducted at Ibn Sina and Khartoum Teaching Hospitals-Otolaryngology Departments (E.N.T), in the period from 1 April – to the 30 of July 2002. Information from both the control and study subjects was taken, using protested questionnaire. Twenty-five normal subjects (50 ears) were taken as a control group selected randomly from the healthy population; their hearing thresholds were tested in order to find a reference level. Seventy-one consecutive patients with uni - or bilateral chronic suppurative otitis media (100 perforated ears) were taken as a study group.

Results: The hearing threshold of the control Sudanese subjects was 17dB. This was obtained by computing the mean for the human speech frequencies (500, 1000, 2000Hz). The lower frequencies were affected regardless of side, site and area of the perforation. **Conductive** hearing loss was the commonest type (79%). The commonest site of the perforations was the **central** site (90%) followed by posteroinferior 3%. The

EDITORIAL

commonest area of tympanic membrane perforations was the subtotal area (54%), followed by large central perforations (25%). The greatest hearing loss was found to be in the posteroinferior and antero-superior perforation of the drum compartment.

Conclusion and recommendations From this study, we can conclude that regardless of the area, the site of tympanic membrane perforation is the most important factor affecting hearing threshold. The posterior central perforation having the greater effect for this reason, any patients with the posteroinferior and antero-superior perforations should have repair of tympanic membrane (myringoplasty).

INTRODUCTION

Tympanic membrane perforation represents a hole in the eardrum, establishing a communication between the middle and external ear. One-third of perforations occur in acute otitis media, and 85% will occur in the anterior inferior quadrant, while the remaining 15% in the posterior superior quadrant. Tympanic membrane perforation usually occurs, due to acute or chronic otitis media, trauma, and neoplasm of the middle ear.¹ Chronic otitis media is usually classified into; tubotympanic disease (perforation of the pars tensa) which presents with central or anterior perforation; and the atticointral disease which most commonly involves the pars flaccida.² The effect of the tympanic membrane perforation on hearing threshold depends on the area and site of the perforation. The area given as the percentage of the ear-drum surface, and the site or location describes the quadrant of the drum affected.¹ For the smallest perforation, the reduction in hearing level was restricted to the lower frequencies. However, as the size of perforation increases a decrease in the high frequency is noticed by Uedo, Nokata and Hoshino.⁴ Usually the larger tympanic membrane perforation, the greater the loss of hearing. The site of the perforation, also affects the degree of hearing loss.⁵ If the perforation is small but located directly over the round window, the loss may even be greater than that due to a large perforation located elsewhere.⁶ Chronic otitis media is one of the commonest otological problems among Sudanese, both adults and children (43%) of ear diseases. They present with perforation of the eardrum, discharging ears and variable degrees of hearing loss as observed by Yagi.⁷ This work is intended to study the effect of the area and site of tympanic membrane perforation on hearing threshold in Sudanese patients.

PATIENTS AND METHODS

This study was carried out at Ibn Sina and Khartoum Teaching Hospitals-Otolaryngology Departments (E.N.T), during the period from first of April to the 30th of July 2002. Data from both the control and study subjects was obtained by a questionnaire and analyzed in master sheets, one for each group.

EDITORIAL

Twenty-five normal subjects (50 ears) were taken as a control group. They were selected randomly from normal healthy subjects with no medical problems or history of otological diseases. Their hearing thresholds were tested in order to find a reference level. This was obtained by computing the mean of the speech frequencies (500, 1000, 2000Hz). Seventy-one consecutive subjects with uni - or bilateral chronic suppurative otitis media (showed 100 perforated tympanic membranes) were taken as the study group. They were the first patients to report to the E.N.T departments at Ibn Sina and Khartoum Teaching Hospitals. Only Sudanese patients without any concomitant medical problems were included. Patient with other ear diseases were excluded e.g past ear surgery, hypertension, diabetes and patients in whom pure tone audiometry was not obtained

Tympanic membrane perforations were classified arbitrary according to site of perforation into six types : antero-superior, antero-inferior ,postero-superior , Postero-inferior ,central ,and attic perforations. Perforations area was classified into four types according to the area of perforation; small ($< \frac{1}{4}$ of tympanic membrane area), medium(up to $\frac{1}{2}$),large (up to $\frac{3}{4}$)and subtotal ($> \frac{3}{4}$ of the). All subjects, both the control and the study groups, were subjected to detailed history to confirm or refute the inclusion or exclusion criteriaand general E.N.T examination.The degree of hearing loss classification was based on the Northern and Downs, classification as: -Normal hearing (0 – 25 dB), mild hearing loss (26 – 40) ,moderate loss (41 – 55), moderate to severe loss (56 – 70), severe loss (71 – 90 dB) and(>91 dB H.L)as profound loss .⁸

RESULTS

Control group: Includes 12 males and 13 females with a mean age of 33.5 years. The average pure tone threshold of both was 17 dB for air conduction, and 10 dB for bone conduction.

Study group: There were (71) patients, (29 male and 42 females with a mean age of 26 years .These patients had (100) perforated ears, (29) patients showed bilateral perforations, (22) showed right-sided perforations and (20) patients showed left sided perforations. Table1and Table2 show the different sites and areas of perforations and the mean air conduction and bone conduction losses.

Most of the patients (79%) presented with conductive hearing loss (15%) with a mixed type of hearing loss and (6%) showed normal hearing thresholds. ⁸

EDITORIAL

The most affected area was central (90%) followed by Postero-inferior perforations 3%. (Table1) most of the patients presented with subtotal perforations (54%) followed by the large tympanic membranes perforations (25%). (Table2)

The most severe effect of site on hearing thresholds was encountered with Postero-inferior perforations and antero-superior perforations.

DISCUSSION

The control group, both males and females showed an average air conduction threshold of 17dB (HTL) which is within the international reference range.⁸ In this study group, the age and sex of the patients were found to have no effect on hearing threshold, irrespective to the site and size of perforation. This in agreement with those of **Yung**.⁹ The low frequencies were noted to be affected first, and with a higher threshold shift than the mid and high frequencies. This is in accord with **Bamanie** who did a similar study in Saudi patients and American Academy of Otolaryngology.^{10, 11} As the area of the perforation increases the hearing threshold increases, with marked affection at the mid and higher frequencies, and a bigger air-bone gap becomes apparent. Also this finding agrees with **Ueda, Nokata and Hoshino**.⁴ In this study, it was found that the effect of the site of tympanic membrane perforation on hearing threshold was more marked when the perforation was in the postero-inferior quadrant compared to other sites which is similar to **Austin**.^{12,13} It is to be recalled that ninety percent (90%) of the perforations in this study were central with minimum effects on hearing thresholds. The postero-inferior small perforations have greater effect on hearing thresholds due to the loss of the baffle effect. This is similar to **Madigan Army Medical Centre**.¹

EDITORIAL

CONCLUSIONS AND RECOMMENDATIONS

The hearing threshold for both males and females in the control Sudanese subjects was 17dB (HTL). The age and sex of the patients have no effect on the site and area of the tympanic membrane perforation. The commonest site was the central perforation. The commonest area was subtotal perforation. Conductive hearing loss was the commonest type of hearing loss(79%,).

The site of tympanic membrane perforations was found to be more important than the area affected. Small postero-inferior and antero-superior perforations have the highest mean air conduction loss of 50- 54 dB.

We recommend that patients with posterior-inferior and antero-superior perforations should have repair of tympanic membrane (myringoplasty) to restore hearing.

ACKNOWLEDGEMENTS

The authors would like to thank Professor Osman Mohamed Elmustafa for his alternations in this article and his invaluable comments.

REFERENCES

- 1- Madigan Army Medical Center Tympanic membrane perforation: Referral guideline 2002; www.mamc.amedd.army.mil.
- 2- Berger G. Nature of spontaneous tympanic membrane perforation in acute otitis media in children. *Journal of Laryngology and Otology*1989; 103: 1150-1153.
- 3- Mills R.P Management of chronic suppurative otitis media. Scott Brown's Otolaryngology, 6th ed. Alan G Kerr (General Editor). Butterworth-Heinemann (Pub) 1997; Vol. 3(10): 1-15.
- 4- Ueda, H., Nakata-S, and Hoshino, M. The effects of Effusion in the Middle Ear and the 5- Perforation of Tympanic Membrane on Otoacoustic Emissions in Guinea Pigs. *Otolaryngology Department, Nagoya University* 1998; 122(1-2): 41-46
- 5- American Academy of Otolaryngology–Head and Neck Surgery www.intrnet-org/ENT Net health into/ear/perforation 2002;**13**/04/23.

EDITORIAL

- 6- Gina berg, Irwin and White-Thomas P Otological disease and examination. In: Katz's (Ed.) Handbook of clinical audio logy, 4th Edition Baltimore, Williams & Wilkins. 1994; P.14.
- 7-Yagi, H.I, The Pattern of Diseases of the Ear in Sudanese Patients. Sudan Medical Journal, Sudan Medical Association. 1989; Vol. 27 (1-4): 44-47.
- 8-Ross J.M, Rosservalents, L. and Holly-Hosford Dunn Application in Diagnostic Audiology. Audiology Diagnosis. Andreoseils (Ed). Thieme (Pub) New York. 2000; 11, 227-251.
- 9-Yung M.W. Myringoplasty: Hearing Gain in Relation to Perforation Site. The Journal of Otolaryngology 1983; 97: 11-17.
- 10- Bamanie A.H. Effect of tympanic membrane perforation on hearing loss among Saudi adults, a thesis submitted for MSC Degree in Audio logy. 1996; 1-144
- 11- American Academy of Otolaryngology –Head and Neck Surgery Acoustic effects of middle ear-disease. Acoustical aspects of chronic ear surgery. Goode R.L (Ed). American Academy of Otolaryngology 1987; (Pub) 80187: 27-28.
- 12- Austin D.F. Sound conduction of the diseased ear . Journal of Laryngology and Otology 1978; 92/5, 367-393.
- 13- Anthony W.P., Harrison C.M., Tympanic membrane perforation: effect on audiogram. Archives of Otolaryngology 1972 ; 95: 506-510.

EDITORIAL

<i>Site of perforation</i>	<i>No. of patients</i>	<i>Percent</i>	<i>Mean air conduction(dB)</i>	<i>Mean bone conduction(dB)</i>
<i>Antero-superior</i>	2	2%	50	16
<i>Antero -inferior</i>	2	2%	45	12
<i>Postero- superior</i>	2	2%	48	13
<i>Postero- inferior</i>	3	3%	54	14
<i>Central</i>	90	90%	37	11
<i>Attic</i>	1	1%	24	15
<i>Total</i>	100	100%	–	–

*Table1:
Distribution of site of perforation among the study group*

Table 2: Distribution of area of perforation among the study group

<i>Size of perforation</i>	<i>No. of patients</i>	<i>Percent</i>	<i>Mean air conduction (dB)</i>	<i>Mean bone conduction (dB)</i>
<i>Small</i>	9	9%	23.7	0
<i>Medium</i>	12	12%	30.0	-2
<i>Large</i>	25	25%	41.0	1
<i>Subtotal</i>	54	54%	39.7	2
<i>Total</i>	100	100%	–	–