

**POLYMORPHISMS OF PFCRT AND PFMDR-1 GENES AND CHLOROQUINE RESISTANCE OF *P. FALCIPARUM* IN WAD MEDANI (CENTRAL SUDAN)**

**Mohammed Osman A/Wahed Madani<sup>1\*</sup> Ahmed El Tahir<sup>2</sup>, Adil Mergani<sup>1</sup>, Mawia Al Balal<sup>2</sup>, Mohammed Elamien<sup>2</sup> & Nasr Eldin. Elwali<sup>1</sup>**

1. Institute of Nuclear Medicine, Molecular Biology & Oncology , University of Gezira

2. Faculty of Medicine, University of Gezira-Sudan

\* **Corresponding author:** Mohammed Osman Abd El Wahed Madani: *Institute of Nuclear Medicine, Molecular Biology & Oncology, University of Gezira, Sudan.* Fax: 249-511-46640 e-mail: mohammed\_abdelwahid@yahoo.com

**ABSTRACT**

**Introduction:** Malaria parasite resistant to Chloroquine poses severe and increasing health problems in tropical countries. Monitoring the drug resistance by implementing the molecular markers may be essential to overcome the problem, therefore this study aims to assess the Chloroquine resistance of Plasmodium Falciparum parasite in central Sudan, using molecular markers.

**Methods:** One hundred and seventy six patients were confirmed *P. falciparum* positive. Sixty-four were selected and only forty patients completed the follow-up. In vivo sensitivity assay was used accompanied with standard regimen of Chloroquine phosphate. DNA was extracted from blood on filter paper (day 0) and was used to amplify two genes *P. Falciparum* transporters gene *Pfcrt* and multi-drug resistant gene-1 *Pfmdr-1*.

**Results:** Among forty patients, 54% responded to Chloroquine regimen with adequate clinical response (ACR), however, 46% showed treatment failure. All treatment failures were treated with Artemether or Quinine. The amplification of *Pfcrt* gene (n, 18) and *Pfmdr 1* gene (n, 29), had shown that 72% of *Pfcrt* T76 were mutant allele, 22% were K76 wild-type, however, only 5% were mixed alleles T/K. while *Pfmdr 1* gene (n, 29) revealed that 55% were wild genotype N 86, 38% were mutant Y 86, and 7% were mixed alleles Y/ N 86.

**Conclusion:** The high frequency of the mutant *Pfcrt* 76T gene among *P.*

*Falciparum* isolates was consistent with in vivo study supports the hypothesis that *Pfcrt* 76T gene could be used as predictive marker for Chloroquine susceptibility in epidemiological surveys.

**Key words:** *P. Falciparum*, drug resistance, *in vivo*, PFCRT, PFMDR 1and RFLP

تشكل الملاريا المقاومة للكلوروكوين خطراً متنامياً في الدول المدارية، لذلك فإن استخدام (الموسمات الجزيئية) في استقصاء مقاومة الطفيل لعقار الكلوروكوين قد تشكل أهمية في التغلب على هذه المشكلة. هدفت هذه الدراسة لاستقصاء مقاومة الطفيل (بلازموديوم فالسبارم) لعقار الكلوروكوين في وسط السودان باستخدام الموسمات الجزيئية. شملت هذه الدراسة مائة ستة وسبعون تأكيد أنهم موجوبون للفحص، تم اختيار أربعة وستون منهم لهذه الدراسة، فقط أربعون تمت متابعتهم كاملاً. استخدمت طريقة متابعة حساسية الطفيل في داخل المريض طبقاً للجرعة القياسية لفوسفات الكلوروكوين وتم استخلاص الحمض النووي الرايبوزي منقوص الأكسجين (DNA) من ورقة ترشيح بها دم أخذ قبل بدء العلاج لكل مريض. واستخدمت لتقييم وتحديد مدى انتشار الطفرات المقاومة للكلوروكوين في اثنين من مورثات طفيل البلازموديوم فالسبارم (PFCRT K76T و PFMDR.1 N86Y) باستخدام جهاز التفاعل التسلسلي والإنزيمات القاطعة للحمض النووي. شملت هذه الدراسة أربعون مريضاً، استجاب 45% منهم للعلاج بينما لم يستجب 46% منهم للعلاج وتم علاج كل المرضى الذين لم يستجيبوا للعلاج بعقار الارتيميثر أو الكينين. أجري المسح الوراثي للطفرات باستخدام تفاعل التسلسلي البلمر (PCR) والإنزيمات القاطعة للحمض النووي (RFLP) للمورثين (PfCRT) عددها (18) و (Pfmdr-1) عددها (29) ووجد أن 72% من جين (PfCRT) تحمل طفرات (T76) للجين (PfCRT) وأن 22% بالمورث (K76 PfCRT) بينما وجد أن 6% تحمل طفرات خليط من الأليل الطافر وغير الطافر (K/T76) أو وضحت نتيجة المسح الوراثي للجين (Pfmdr-1 Y86N) بأن 38% تحمل طفرات وأن 55% من الطفيليات لا تحمل طفرة له وأن 7% من الطفيليات تحمل خليط من الأليل الطافر وغير الطافر. خلصت الدراسة أن زيادة تكرار الجين الطافر (PfCRT) في طفيليات البلازموديوم فالسبارم تعزز افتراضية إمكانية استخدام جين (PfCRT 76T) موسم أساسي لمراقبة ومعرفة حساسية مقاومة طفيل البلازموديوم فالسبارم لعقار الكلوروكوين في المسوحات الوبائية.

## INTRODUCTION

Resistance to Chloroquine was noted in Thai-Cambodian border and in Colombia since the late 1950s (1), whereas in Africa, the chloroquine resistance was first reported in east of Africa in 1978 and the resistance spread to centre and south of the continent and at last to the west of Africa 1983, and by 1989 chloroquine resistance was wide spread in Sub-Saharan Africa (2).

Polymorphism in two genes of Plasmodium falciparum (P.falciparum) genome had contributed in chloroquine resistance. The P. facibarum chloroquine resistance transporters (PfCRT) gene located in chromosome 7 and this segment harbors a 13-exon, PfCRT, having point mutations that associate completely with chloroquine resistant (CQR) in parasite lines from Asia, Africa, and South America (3),

Laboratory studies in Cameroon and Mali have strongly suggested that P. falciparum chloroquine resistance transporter (PfCRT Lys76Thr) gene can be used as a marker in surveillance for chloroquine-resistant falciparum malaria as a molecular marker (4-6). In diverse geographical areas several studies support the association between the PfCRT K76T and chloroquine resistance in Africa (Mali) (5), Sudan (7-9), Nigeria (10), Gabon (11), Gambian (12), Malawi (13), Liberia (14), Tanzania (15), Mozambique (16), Ghana (17), Camroon (18) and Senegal (19-20). Asia Thailand (21) and India (22), and Amazon region Brazil (23, 24), however studies from Uganda at the south gate of Sudan reported no association (25).

Initial studies of the gene P. falciparum Multidrug resistance gene (Pfmdr-1) located on chromosome 5, illustrate that an Asn→Tyr mutation at amino acid 86 (N86Y) and other mutations in this gene correlated with chloroquine resistance in {Sudan (7), Mali (5), Gambia (12), Uganda (27), Brazil and Thailand (21)}, however, some authors from {Uganda (28), Brazil (29), and Thailand (30)}, could not find this correlations in their studies.

## MATERIAL AND METHODS

**Patients:** This study was conducted in Marengan Health Center and 48 patients with positive P. Falciparum bloods were enrolled in the study.

### The In vivo study:

Chloroquine phosphate (AMIPHARMA Laboratories Ltd. Sudan B N/T0013) was administrated orally at a dose of 10 mg per kilogram (kg) of body weight in day (0) and 5 mg per kg body weight after 6 hours, and in day 1 and day 2, the patients were observed after each dose, and they were given another full dose or a half dose if vomiting occurred within 30 min or within 31 min to 1 hour respectively. Clinical and

## **EDITORIAL**

parasitological follow-up was done on day 1, 3, 7 and 14. Whenever symptoms were reported or parasiteima increased, the case is considered as in vivo Chloroquine resistance state (31) and alternative treatment was scheduled for that.

### **Assessment of in vivo response and implementing molecular techniques:**

This was done according to WHO assessment (31). Blood samples were obtained on white man filter paper 0.3 immediately before treatment and stored in 4°C, DNA was extracted from filter paper blood samples by cutting an approximately 4 mm<sup>2</sup> piece of paper was placing in 100 µl methanol, air dried, and incubated at 95 to 100 °C in 50 µl of water for 15 minutes. During incubation, the tube was subjected to high-speed vortex three times. 5 µl of resulting solution used as template for PCR (5)

The PCR reaction was performed in 30 µl volumes which contains 1X PCR Buffer II (Gene Amp® 10X Buffer II [100 mM Tris-HCl pH 8.3, 500 mM KCl], Applied Bio system), 1.5 mM MgCL<sub>2</sub> (MgCl<sub>2</sub>, Applied Bio system), 0.2 mM each of the dNTP (Gene Amp® dNTPs, Applied Bio system), 1.0 µ mol of each sense and antisense primers 1 U of AmpliTag Gold (Applied Bio system), 5 µl of the Template DNA; and the reaction volume was completed to 30 µl by ddH<sub>2</sub>O.

Nested PCR for the detection of tyr-86 allele of pfmdr-1 in field isolates for all reactions was amplified using Gene Amp® PCR system 9700 (Applied Bio system) in the following conditions: one cycle at 94 °C for 3 minutes; 40 cycles at 94 °C for 1 minute, 49 °C for 1 minute, and 72° C for 1 minute; and a final extension at 72° C for 10 minutes was carried out using outer primers MdrA1 and MdrA3 (32). 3 µl were used for the second round as a template of nested PCR, with a new set of inner primers Mdr A2 and Mdr A4 without change in the PCR conditions as described by (32). A nested PCR protocols were used to identify the K76T in the Pfcrf gene (3) with a slight modification on primers used to amplify in Gene Amp® PCR system 9700 (Applied Bio system). The outer primers tcrp- 1 and Pfcrf-13 in the following conditions: one cycle at 94° C for 3 minutes; 40 cycles at 94 °C for 30 seconds, 55 °C for 30 seconds, and 60° C for 30 seconds; and a final extension at 60 °C for 3 minutes. 2 µl of product was diluted 1/20 and 3 µl were used as a template for the nested reaction using Pfcrf-14 and Pfcrf-15 primers in the following conditions: one cycle at 94 °C for 3 minutes; 40 cycles at 92 C for 30 seconds, 49 °C for 30 seconds, and 65 °C for 30 seconds; and a final extension at 65 °C for 3 minutes. For each series of samples, water was used as a negative control, HB3-strain DNA was used as the wild-type control, and Dd2 DNA was used as the mutant control. All the amplifications were checked using agarose gel electrophoresis, in which PCR products were run in 1.5 % ( Pfmdr-1 gene) and 2 % ( Pfcrf gene).

Comparing gene sequence in group of individuals; change in one or more base pairs occur causing variation in the sequence, these variations are called polymorphisms, furthermore, restriction site may be created or abolished at that point, this is known as Restriction Fragment Length Polymorphism (RFLP) (33) . The mutations in the two genes Pfcrf and Pfmdr-1 have been abolished by the restriction site for Apo I enzyme (Fig 2.4). Digestion with this enzyme was used for typing these polymorphisms. In a total volume of 15 µl, 2 µl PCR product were digested overnight at 50°C with 1 U Apo I, 1.5 µl 10x NE Buffer 3 (100 mM NaCl, 50 mM Tris-HCl, 10 mM magnesium chloride and 1mM dithiothreitol (pH 7.9 @ 25 °C), 0.15 µl of 100X BSA (200µg/ml ) and deionized water. Using the Genomic DNA from strain 3D7 and Dd2 were Amplified and digested in the same way serving as control for complete digestion, and undigested fragment respectively (Fig. 2.1).

EDITORIAL




Figure 1 Recognition site for Apo I enzyme

**Statistical analysis:** SPSS (Statistical Package for Social Science) software was used for statistical analysis.

**RESULTS**

**Study area:** The study was conducted at Marengan Health Center Medani, Sudan, from September to December 2002, where the malaria incidence is high.

**Patients:** Patients with symptoms or signs suggesting malaria were screened for the presence of malaria parasites in their peripheral blood. One hundred and seventy six (176) patients were found to be smear positive for malaria; 94% of them were infected with *P. falciparum*; 6 % with mixed infection *P. falciparum* and *P. vivax* or *P. vivax* only.

Sixty four (64) patients fulfilled the criteria for selection and enrolled in the study; and only 40 patients were completed the period of follow up.

**Response of *P. falciparum* isolates to chloroquine according to in vivo test:**

Fig (2) shows the adjustment of isolates to chloroquine according to WHO standard regimen dose and selection criteria. From forty (40) patients treated with chloroquine, only 22 (54%) responded fully with adequate clinical and parasitological response (ACR), in 18 (46%) patients drug resistance was found distributed to 9 (23%) with early treatment failure (ETF) and 9 (23%) with late treatment clinical and parasitological failure (LTF). All the treatment failures were treated with artemether injection 1.6 mg/ Kg body weight or quinine and followed as required, there was no recrudescence reported.




Figure 2: The distribution *P. falciparum* isolates according to in vivo sensitivity to chloroquine.

Table 1 shows the distribution of the mean parasitemia and the standard error of in vivo test. According to sensitivity all the 22 (54%), the group of ACR, with mean initial parasitemia,  $20089 \pm 6520$ ,  $5723 \pm 2713$ , and  $00 \pm 00$  in D0, D1, and D3 respectively. Finally, only one patient was found to have severe malaria

**EDITORIAL**

symptoms before day 3, and treated with quinine as alternative drug and followed as required.


**Table 1:** The parasitaemia mean ± standar error of the mean (S. E. M) according to in vivo sensitivity response to chloroquine

<i>In vivo</i> Sensitivity		Number	Mean ± (S. E. M)
D0	ACR	22	20089 ± 6520
	ETF	9	22550 ± 6532
	LTF	9	16230 ± 7083
D1	ACR	22	5723 ± 2713
	ETF	9	3530± 1180
	LTF	9	5388 ± 3934
D3	ACR	22	00 ± 00
	ETF	8	1869 ± 1093
	LTF	9	00 ± 00
D7	ACR	22	00 ± 00
	ETF	-	-
	LTF	8	1015 ± 541
D14	ACR	22	00 ± 00
	ETF	-	-
	LTF	1	400 ± 00

**The prevalence of PfCRT T76 and Pfmdr 1 mutations:**

Fig. (3) Presented the pattern of detection of *P. falciparum* chloroquine resistance transporter gene (PfCRT) using nested PCR method where as, (Fig. 4) shows the detection of *P. falciparum* multi-drug resistance 1 gene using nested PCR method. (Fig. 5), represented the pattern of restriction fragment length polymorphism (RFLP) of Pfmdr 1 gene. The prevalence of PfCRT T76 and Pfmdr 1 Y86 was screened in 18 and 29 parasite isolates respectively. Restriction fragment length polymorphism (RFLP) technique using Apo 1 enzyme was used for screening the isolates. Table 2 shows that the PfCRT T76 mutation was present in 13 (72%) samples, 4 (23%) had the wild-type of PfCRT K76 and only one (5%) sample with mixed alleles of wild and mutant (T76K). and the prevalence of Pfmdr-1 Y86 was present in 11 (38%) mutant type and 16 (55%) were found to be Pfmdr 1 N86 and 2 (7%) carried the mixed alleles of wild and mutant type (Y86N) in (Table 3.).


**Figure.3.:** Detection of of *P. falciparum* chloroquine resistance transporter gene (PfCRT) using nested PCR method.



**EDITORIAL**


**Fig. 4:** Detection of of *P. falciparum* multi-drug resistance 1 gene using nested PCR method.

Lane 1: M: 100-bp DNA marker. Lane 2: PCR product (positive control). Lanes 3 -11 Positive amplification (502 bp) PCR products and lane 12 negative control.




**Figure.5:** PCR-RFLP screening of *P. falciparum* isolate for codon 76 polymorphism. Lane 1 and 10: DNA markers.

Lane 2: digested PCR product (positive control). Lane 3&5: K76 polymorphism (cutted by Apo1 restriction enzyme to 145 and 64 bp). Lane 4, 6, 7, 8&9: T76 polymorphism (uncutted full length 209 Pb).



**Figure.6:** PCR-RFLP screening of *P. falciparum* isolate for codon 86 polymorphism

Lane 1: Mixed infection by two strains carrying both haploid alleles N86 and Y86 Lane 2, 4 & 7: N86 allele (cutted by Apo1 restriction enzyme to 254 and 248 bp) Lanes 3, 5, 6 & 8: Y86 allele (uncutted by Apo1 restriction enzyme full length 502 bp). Lane 9: undigested PCR product (positive control). Lane 10: DNA marker.



**Association of mutations and treatment outcome:**

In (Table 2) six samples were classified as ACR according to in vivo sensitivity test, the molecular technique shows three samples had K76, two of samples had T76 (the mutant type), and the last one with mix alleles T76 and K76. In eight isolates classified as ETF using in vivo test, the molecular techniques showed only one sample (12.5%) which had the wild type K76 and the rest of the isolates (87.5%) carrying

**EDITORIAL**

the mutant one T 76. In the isolates classified as LTF all isolates (100%) had the mutant type T76. No sample in treatment failure outcome of Pfcr1 gene was observed to have mixed alleles.

Pfmdr 1 gene mutation represented in (Table. 3) no sample in the group, which classified as ACR of Pfmdr-1 gene was observed to have mixed alleles and only two isolates with mixed alleles in isolate classified as ETF were found.

**Frequency of Pfcr1 and Pfmdr 1 genotypes among P. falciparum isolates:**

The distribution P. falciparum isolates according to Pfmdr-1 and Pfcr1 genotypes were illustrated in (Fig.7)

**Table 2:** Distribution of Pfcr1 genotypes in P.Falciparum isolates according to in vivo sensitivity to chloroquine

Total	Pf Mdr-1 marker			In vivo sensitivity
	N/Y	Y	N	
14	0	4	10	ACR
9	2	3	4	ETF
6	0	4	2	LTF
29	2	11	16	Total

Pfcr1  $P = 0.002$  in vivo sensitivity  $P = 0.000$

**Table 3:** Distribution of Pfmdr-1 genotypes in P.falciparum isolates according to in vivo sensitivity to chloroquine

Total	Pfcr1 markers			In vivo sensitivity
	K/T	T	K	
6	1	2	3	ACR
8	0	7	1	ETF
4	0	4	0	LTF
<b>18</b>	<b>1</b>	<b>13</b>	<b>4</b>	<b>Total</b>

Pf Mdr-1 marker 1  $P = 0.005$

In vivo sensitivity  $P = 0.000$

**EDITORIAL**

**Table 4:** In vivo, and markers marker results for Pfprt gene and Pfmdr-1 genes in all isolates

Isolate	<i>In vivo</i>	Pfmdr 1	Pfprt
C10	ACR	ND	ND
C11	ACR	ND	ND
C12	ETF	Y	T
C13	ACR	N	K
C15	ACR	N	ND
C16	ETF	N	T
C19	ACR	N	ND
C20	ACR	N	ND
C21	LTF	ND	ND
C22	ETF	ND	ND
C23	ETF	Y	T
C24	ACR	N	K
C25	ACR	Y	ND
C27	ACR	N	ND
C28	ACR	N	ND
C29	LTF	N	T
C30	LTF	Y	T
C31	ACR	ND	ND
C33	ETF	Y	ND
C34	ACR	ND	ND
C35	ACR	ND	ND
C36	ACR	ND	ND
C37	ACR	ND	ND
C38	ETF	N	K
C40	ETF	N/Y	T
C41	ACR	Y	ND
C42	ETF	N	T
C43	ACR	Y	T
C44	ACR	N	ND
C46	LTF	Y	T
C47	LTF	Y	ND
C48	ETF	N/Y	T
C49	LTF	N	T
C50	ACR	N/Y	T/K
C51	LTF	N	ND
C/2R	LTF	ND	ND
C5/R	LTF	Y	T
C52	ACR	N	K
C53	ACR	Y	T
C1/R	ACR	ND	ND
Total	40	29	18

**DISCUSSION**

**In vivo assessment:**

In this study, the failure rate for treatment during in vivo assessment of sensitivity to chloroquine was 46%, this is very remarkable because the similar failure rate was reported in Sudan (34), and also similar Frequencies (39-46%) were observed in other African countries (35-38) .

According to the recent classification of WHO criteria for in vivo drug sensitivity into ACR, ETF and LTF (31) , the results distributed as follows; 22 patients (54%) were ACR, 9 patients (23%) ETF and 9 patients were LTF (23%). (Fig. 2).

In (ACR) group, the isolates with low or moderate levels of resistance, the infection could be resolved and cured due to many factors that could be summarized in a proper drug dose, parasites genetic material and partial immunity experiences, this more evident in full immune response patients in endemic areas (39). In case of the (ETF) and (LTF) there are many factors contributing to the treatment outcome; factors that influencing host and parasites; in case of host the main factors are immune status of the patients, for the

## **EDITORIAL**

parasites the mutations, which affect drug action, the interaction between sexual stage and the vectors and moreover, the environment may alter in this process (1, 40-42) , all these factors may allow the parasites to survive in the presence of chloroquine.

In general, residents of malaria-endemic areas sometimes spontaneously clear *P. falciparum* infection without drug treatment, implying an important role for host factors, such as immunity, in this clearance (43).

### **The effect of the initial parasitemia on the treatment outcome:**

Comparison of the effect of the initial parasitemia on the treatment outcome showed no significant difference in chloroquine response at different levels of the initial parasitemia (Table 1), this indicates that; the initial parasitemia has no role in determining the chloroquine in vivo response. This finding was in line with study done in Sennar (44) , however, study done by Wernsdorfer et al, observed higher mean initial parasitemia in the group of resistance parasites, than in sensitive one (45).

### **Molecular markers:**

#### ***P. falciparum* Chloroquine resistance transporter (Pfcr1):**

Characterization of genes controlling drug resistance in parasite, as in dihydrofolates reductase gene lead to the development of molecular markers, this is used for proper diagnosis and early detection of treatment failure (46).

The prevalence of Pfcr1 T76 mutations (the mutations which was incriminated in chloroquine resistance) was present in 78% of our isolates, and 22% (4 isolates) carrying the wild type of Pfcr1 K76 (Fig. 7), most of the isolates with mutation type might have been resistant in vivo, however, 34% and 14% of the mutant alleles T76 and T/K76 were recovered by immune response and classified as ACR ( Table 2), this find indicated the important role of the host immunity, this observation was consistent with a number of studies (28, 43, 47).

Most of the isolates classified as in vivo resistance, carrying Pfcr1 T76, so based on this above observation; we strongly associated the subsequent resistance to chloroquine in vivo with the presence of Pfcr1 T76, at the time of treatment. These data, are in line with genetic evidence of Fidock et al (3), on the other hand, only one isolate was detected carrying the wild type K76, and classified as in vivo resistance (Table 2). The possible explanation is that the patient had mixed infections with low level of resistant parasites carrying the mutant type, under the drug pressure the sensitive parasites were eliminated and the resistant ones expanded resulting in treatment failure. The presence of Pfcr1 T76 at the time of treatment was also strongly associated with subsequent resistance to chloroquine in vivo.

#### ***P. Falciparum* multi drug resistance-1 gene (Pfmdr-1):**

The prevalence of the wild type of Pfmdr-1 isolates is more, compared to the mutant type in the screened samples (Fig. 3), indicating the less sensitivity of Pfmdr-1 for detecting the resistance isolates predicted by in vivo test.

Among ACR group, the N86 can predict 71% of the sensitivity in vivo (Table 3), in the resistance group 50% of isolates carrying the mutant type Y86. This observation is consistent with some other studies in which decrease in association between Pfmdr- 1 Y86 and sub-mutations and chloroquine resistance in vivo was reported (48, 49)

Recently, many investigators showed that resistant infections in vivo can be due to parasites with no Pfmdr-1 mutations at position Y86 and gave strong evidence to the link of this resistance to other gene (3, 5, 10). Chloroquine resistance in Sudan was first reported In the Gezira area (50), and then in Sennar (51) and Gadarif (52), but now resistance to chloroquine was reported in different parts of the country. In the last decade progress has been done in the molecular basis of drug resistance in *P. falciparum*. As more information on the genetics of anti-malarial resistance becomes available, designing of new molecular-based tools for drug resistance evaluation, and applying them in early detection of drugs resistant will help in proper selection of antimalarial drugs and rapid treatment of patients which is the corner stones in malaria control.

**EDITORIAL**

**ACKNOWLEDGEMENT**

I would like to express my deepest thanks to all who participated in this study namely, Dr. Mohamed El Emam; Professor Ali Habour, (Faculty of Medicine, Department of Pediatrics); Dr. Badria Babiker and Professor Ahmed Babiker; Mr. Salah; Mrs. Nahla; Mr. Tilal Babiker Ajeeb (Institute of Tropical Medicine, Ministry of Science and Technology); Dr. Amar Khamis (University of Sudan); Dr. Fathi Al Rabaa (Faculty of Science, University of Khartoum); Husham Bakhet (Faculty of Agriculture, University of Khartoum); Mr. Badr El Deen El Fadil (Faculty of Medical Laboratory Science University of Gezira); Mr. Ismeel Adam, Mr. Al Fatih Ibrahim; Mr. Jalal. H. AL Awad; Mr. Abbas M. El seed and Dr. El Saidig (Marengan Clinical Center, Ministry of Health) .

**REFERENCES**

1. Wernsdorfer, W.H. and D. Payne, *The dynamics of drug resistance in Plasmodium falciparum*. Pharmacol Ther, 1991. 50(1): p. 95-121.
2. Wongsrichanalai, C., A.L. Pickard, W.H. Wernsdorfer, and S.R. Meshnick, *Epidemiology of drug-resistant malaria*. Lancet Infect Dis, 2002. 2(4): p. 209-18.
3. Fidock, D.A., T. Nomura, A.K. Talley, R.A. Cooper, S.M. Dzekunov, M.T. Ferdig, L.M. Ursos, A.B. Sidhu, B. Naude, K.W. Deitsch, X.Z. Su, J.C. Wootton, P.D. Roepe, and T.E. Wellems, *Mutations in the P. falciparum digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance*. Mol Cell, 2000. 6(4): p. 861-71.
4. Basco, L.K., M. Ndounga, V.F. Ngane, and G. Soula, *Molecular epidemiology of malaria in Cameroon. XIV. Plasmodium falciparum chloroquine resistance transporter (PFCRT) gene sequences of isolates before and after chloroquine treatment*. Am J Trop Med Hyg, 2002. 67(4): p. 392-5.
5. Djimde, A., O.K. Doumbo, J.F. Cortese, K. Kayentao, S. Doumbo, Y. Diourte, A. Dicko, X.Z. Su, T. Nomura, D.A. Fidock, T.E. Wellems, C.V. Plowe, and D. Coulibaly, *A molecular marker for chloroquine-resistant falciparum malaria*. N Engl J Med, 2001. 344(4): p. 257-63.
6. Durand, R., V. Huart, S. Jafari, and J. Le Bras, *Rapid detection of a molecular marker for chloroquine-resistant falciparum malaria*. Antimicrob Agents Chemother, 2002. 46(8): p. 2684-6.
7. Babiker, H.A., S.J. Pringle, A. Abdel-Muhsin, M. Mackinnon, P. Hunt, and D. Walliker, *High-level chloroquine resistance in Sudanese isolates of Plasmodium falciparum is associated with mutations in the chloroquine resistance transporter gene pfcr1 and the multidrug resistance Gene pfmdr1*. J Infect Dis, 2001. 183(10): p. 1535-8.
8. Abdel-Muhsin, A.A., M.J. Mackinnon, P. Awadalla, E. Ali, S. Suleiman, S. Ahmed, D. Walliker, and H.A. Babiker, *Local differentiation in Plasmodium falciparum drug resistance genes in Sudan*. Parasitology, 2003. 126(Pt 5): p. 391-400.
9. Ochong, E.O., I.V. van den Broek, K. Keus, and A. Nzila, *Short report: association between chloroquine and amodiaquine resistance and allelic variation in the Plasmodium falciparum multiple drug resistance 1 gene and the chloroquine resistance transporter gene in isolates from the upper Nile in southern Sudan*. Am J Trop Med Hyg, 2003. 69(2): p. 184-7.
10. Adagut, I.S. and D.C. Warhurst, *Plasmodium falciparum: linkage disequilibrium between loci in chromosomes 7 and 5 and chloroquine selective pressure in Northern Nigeria*. Parasitology, 2001. 123(Pt 3): p. 219-24.
11. Binder, R.K., S. Borrmann, A.A. Adegnik, M.A. Missinou, P.G. Kremsner, and J.F. Kun, *Polymorphisms in the parasite genes for pfcr1 and pfmdr-1 as molecular markers for chloroquine resistance in Plasmodium falciparum in Lambarene, Gabon*. Parasitol Res, 2002. 88(5): p. 475-6.
12. Sutherland, C.J., A. Allouche, J. Curtis, C.J. Drakeley, R. Ord, M. Duraisingh, B.M. Greenwood, M. Pinder, D. Warhurst, and G.A. Targett, *Gambian children successfully treated with chloroquine can harbor and transmit Plasmodium falciparum gametocytes carrying resistance genes*. Am J Trop Med Hyg, 2002. 67(6): p. 578-85.

**EDITORIAL**

13. Kublin, J.G., J.F. Cortese, E.M. Njunju, R.A. Mukadam, J.J. Wirima, P.N. Kazembe, A.A. Djimde, B. Kouriba, T.E. Taylor, and C.V. Plowe, *Reemergence of chloroquine-sensitive Plasmodium falciparum malaria after cessation of chloroquine use in Malawi*. J Infect Dis, 2003. 187(12): p. 1870-5.
14. Checchi, F., R. Durand, S. Balkan, B.T. Vonhm, J.Z. Kollie, P. Biberson, E. Baron, J. Le Bras, and J.P. Guthmann, *High Plasmodium falciparum resistance to chloroquine and sulfadoxine-pyrimethamine in Harper, Liberia: results in vivo and analysis of point mutations*. Trans R Soc Trop Med Hyg, 2002. 96(6): p. 664-9.
15. Schneider, A.G., Z. Premji, I. Felger, T. Smith, S. Abdulla, H.P. Beck, and H. Mshinda, *A point mutation in codon 76 of pfprt of P. falciparum is positively selected for by Chloroquine treatment in Tanzania*. Infect Genet Evol, 2002. 1(3): p. 183-9.
16. Mayor, A.G., X. Gomez-Olive, J.J. Aponte, S. Casimiro, S. Mabunda, M. Dgedge, A. Barreto, and P.L. Alonso, *Prevalence of the K76T mutation in the putative Plasmodium falciparum chloroquine resistance transporter (pfprt) gene and its relation to chloroquine resistance in Mozambique*. J Infect Dis, 2001. 183(9): p. 1413-6.
17. Mockenhaupt, F.P., T.A. Eggelte, H. Till, and U. Bienzle, *Plasmodium falciparum pfprt and pfmdr1 polymorphisms are associated with the pfdhfr N108 pyrimethamine-resistance mutation in isolates from Ghana*. Trop Med Int Health, 2001. 6(10): p. 749-55.
18. Basco, L.K. and P. Ringwald, *Analysis of the key pfprt point mutation and in vitro and in vivo response to chloroquine in Yaounde, Cameroon*. J Infect Dis, 2001. 183(12): p. 1828-31.
19. Daily, J.P., C. Roberts, S.M. Thomas, O. Ndir, T. Dieng, S. Mboup, and D.F. Wirth, *Prevalence of Plasmodium falciparum pfprt polymorphisms and in vitro chloroquine sensitivity in Senegal*. Parasitology, 2003. 126(Pt 5): p. 401-5.
20. Thomas, S.M., O. Ndir, T. Dieng, S. Mboup, D. Wypij, J.H. Maguire, and D.F. Wirth, *In vitro chloroquine susceptibility and PCR analysis of pfprt and pfmdr1 polymorphisms in Plasmodium falciparum isolates from Senegal*. Am J Trop Med Hyg, 2002. 66(5): p. 474-80.
21. Lopes, D., K. Rungsihirunrat, F. Nogueira, A. Seugorn, J.P. Gil, V.E. Do Rosario, and P. Cravo, *Molecular characterisation of drug-resistant Plasmodium falciparum from Thailand*. Malar J, 2002. 1(1): p. 12.
22. Vathsala, P.G., A. Pramanik, S. Dhanasekaran, C.U. Devi, C.R. Pillai, S.K. Subbarao, S.K. Ghosh, S.N. Tiwari, T.S. Sathyanarayan, P.R. Deshpande, G.C. Mishra, M.R. Ranjit, A.P. Dash, P.N. Rangarajan, and G. Padmanaban, *Widespread Occurrence of the Plasmodium Falciparum Chloroquine Resistance Transporter (Pfprt) Gene Haplotype Svmnt in P. Falciparum Malaria in India*. Am J Trop Med Hyg, 2004. 70(3): p. 256-259.
23. Vieira, P.P., M.U. Ferreira, M. Das Gracias Alecrim, W.D. Alecrim, L.H. Da Silva, M.M. Sihuinha, D.A. Joy, J. Mu, X.Z. Su, and M.G. Zalis, *pfprt Polymorphism and the Spread of Chloroquine Resistance in Plasmodium falciparum Populations across the Amazon Basin*. J Infect Dis, 2004. 190(2): p. 417-24.
24. Vieira, P.P., M. das Gracias Alecrim, L.H. da Silva, I. Gonzalez-Jimenez, and M.G. Zalis, *Analysis of the PfCRT K76T mutation in Plasmodium falciparum isolates from the Amazon region of Brazil*. J Infect Dis, 2001. 183(12): p. 1832-3.
25. Talisuna, A.O., J. Kyosiimire-Lugemwa, P. Langi, T.K. Mutabingwa, W. Watkins, E. Van Marck, T. Egwang, and U. D'Alessandro, *Role of the pfprt codon 76 mutation as a molecular marker for population-based surveillance of chloroquine (CQ)-resistant Plasmodium falciparum malaria in Ugandan sentinel sites with high CQ resistance*. Trans R Soc Trop Med Hyg, 2002. 96(5): p. 551-6.
26. Cowman, A.F., S. Karcz, D. Galatis, and J.G. Culvenor, *A P-glycoprotein homologue of Plasmodium falciparum is localized on the digestive vacuole*. J Cell Biol, 1991. 113(5): p. 1033-42.
27. Flueck, T.P., T. Jelinek, A.H. Kilian, I.S. Adagu, G. Kabagambe, F. Sonnenburg, and D.C. Warhurst, *Correlation of in vivo-resistance to chloroquine and allelic polymorphisms in Plasmodium falciparum isolates from Uganda*. Trop Med Int Health, 2000. 5(3): p. 174-8.
28. Dorsey, G., M.R. Kamya, A. Singh, and P.J. Rosenthal, *Polymorphisms in the Plasmodium falciparum pfprt and pfmdr-1 genes and clinical response to chloroquine in Kampala, Uganda*. J Infect Dis, 2001. 183(9): p. 1417-

**EDITORIAL**

20.

29. Pova, M.M., I.S. Adagu, S.G. Oliveira, R.L. Machado, M.A. Miles, and D.C. Warhurst, *Pfmdr1 Asn1042Asp and Asp1246Tyr polymorphisms, thought to be associated with chloroquine resistance, are present in chloroquine-resistant and -sensitive Brazilian field isolates of Plasmodium falciparum*. *Exp Parasitol*, 1998. 88(1): p. 64-8.
30. Chaiyaroj, S.C., A. Buranakiti, P. Angkasekwina, S. Looessuwan, and A.F. Cowman, *Analysis of mefloquine resistance and amplification of pfmdr1 in multidrug-resistant Plasmodium falciparum isolates from Thailand*. *Am J Trop Med Hyg*, 1999. 61(5): p. 780-3.
31. WHO, W.H.O., *Assessment of therapeutic efficacy for uncomplicated falciparum malaria in areas with intense transmission*. Geneva, WHO, 1996. (WHO/MAL/96.1077).
32. Duraisingh, M.T., L.V. von Seidlein, A. Jepson, P. Jones, I. Sambou, M. Pinder, and D.C. Warhurst, *Linkage disequilibrium between two chromosomally distinct loci associated with increased resistance to chloroquine in Plasmodium falciparum*. *Parasitology*, 2000. 121 ( Pt 1): p. 1-7.
33. Winter., P.C., G.I. Hickey., and H.L. Fletcher., *Genetic diseases*, in *Instant notes in Genetics*, B.D. Hames, Editor. 1998, Bios: Leeds. p. 289-292.
34. Yousif, M.A. and A.A. Adeel, *Antimalarials prescribing patterns in Gezira State: precepts and practices*. *East Mediterr Health J*, 2000. 6(5-6): p. 939-47.
35. Yavo, W., E.I. Menan, T.A. Adjetey, P.C. Barro-Kiki, L. Nigue, Y.J. Konan, N.G. Nebavi, and M. Kone, *[In vivo sensitivity of Plasmodium falciparum to amino-4-quinolines and sulfadoxine pyrimethamine in Agou (Ivory Coast)]*. *Pathol Biol (Paris)*, 2002. 50(3): p. 184-8.
36. Villadary, I., C. Paquet, E. Hemelsdael, G. Blanchard, and Z.M. Saki, *[In vivo drug sensitivity of Plasmodium falciparum in the Tabou region of Ivory Coast]*. *Bull Soc Pathol Exot*, 1997. 90(1): p. 10-3.
37. Landgraf, B., H. Kollaritsch, G. Wiedermann, and W.H. Wernsdorfer, *Plasmodium falciparum: susceptibility in vitro and in vivo to chloroquine and sulfadoxine-pyrimethamine in Ghanaian schoolchildren*. *Trans R Soc Trop Med Hyg*, 1994. 88(4): p. 440-2.
38. Rukaria, R.M., S.B. Ojwang, J.B. Oyieke, and C.B. Kigundu, *In vivo and in vitro response of Plasmodium falciparum to chloroquine in pregnant women in Kilifi district, Kenya*. *East Afr Med J*, 1992. 69(6): p. 306-10.
39. Hastings, I.M. and U. D'Alessandro, *Modelling a predictable disaster: the rise and spread of drug-resistant malaria*. *Parasitol Today*, 2000. 16(8): p. 340-7.
40. Wongsrichanalai, C., J. Sirichaisinthop, J.J. Karwacki, K. Congpuong, R.S. Miller, L. Pang, and K. Thimasarn, *Drug resistant malaria on the Thai-Myanmar and Thai-Cambodian borders*. *Southeast Asian J Trop Med Public Health*, 2001. 32(1): p. 41-9.
41. Winstanley, P., *Modern chemotherapeutic options for malaria*. *Lancet Infect Dis*, 2001. 1(4): p. 242-50.
42. Wernsdorfer, W.H., *Epidemiology of drug resistance in malaria*. *Acta Trop*, 1994. 56(2-3): p. 143-56.
43. Djimde, A.A., O.K. Doumbo, O. Traore, A.B. Guindo, K. Kayentao, Y. Diourte, S. Niare-Doumbo, D. Coulibaly, A.K. Kone, Y. Cissoko, M. Tekete, B. Fofana, A. Dicko, D.A. Diallo, T.E. Wellems, D. Kwiatkowski, and C.V. Plowe, *Clearance of drug-resistant parasites as a model for protective immunity in Plasmodium falciparum malaria*. *Am J Trop Med Hyg*, 2003. 69(5): p. 558-63.
44. Kouznetsov, R.L., W. Rooney, W.H. Wernsdorfer, A.A. El Gaddal, D. Payne, and R.E. Abdalla, *Use of the in vitro microtechnique for the assessment of drug sensitivity of Plasmodium falciparum in Sennar, Sudan*. *Bull World Health Organ*, 1980. 58(5): p. 785-9.
45. Wernsdorfer, W.H., B. Landgraf, G. Wiedermann, and H. Kollaritsch, *Chloroquine resistance of Plasmodium falciparum: a biological advantage?* *Trans R Soc Trop Med Hyg*, 1995. 89(1): p. 90-1.
46. Cortese, J.F. and C.V. Plowe, *Antifolate resistance due to new and known Plasmodium falciparum dihydrofolate reductase mutations expressed in yeast*. *Mol Biochem Parasitol*, 1998. 94(2): p. 205-14.
47. Pillai, D.R., A.C. Labbe, V. Vanisaveth, B. Hongvangthong, S. Pomphida, S. Inkathone, K. Zhong, and K.C. Kain, *Plasmodium falciparum malaria in Laos: chloroquine treatment outcome and predictive value of*

**EDITORIAL**

- molecular markers*. J Infect Dis, 2001. 183(5): p. 789-95.
48. Basco, L.K. and P. Ringwald, *Molecular epidemiology of malaria in Yaounde, Cameroon. III. Analysis of chloroquine resistance and point mutations in the multidrug resistance 1 (pfmdr 1) gene of Plasmodium falciparum*. Am J Trop Med Hyg, 1998. 59(4): p. 577-81.
49. McCutcheon, K.R., J.A. Freese, J.A. Freaan, B.L. Sharp, and M.B. Markus, *Two mutations in the multidrug-resistance gene homologue of Plasmodium falciparum, pfmdr1, are not useful predictors of in-vivo or in-vitro chloroquine resistance in southern Africa*. Trans R Soc Trop Med Hyg, 1999. 93(3): p. 300-2.
50. Omer, A.H., *Response of Plasmodium falciparum in Sudan to oral chloroquine*. Am J Trop Med Hyg, 1978. 27(5): p. 853-7.
51. Ibrahim, M.E., F.M. Awad-el-Kariem, I.M. el Hassan, and E.R. el Mubarak, *A case of Plasmodium falciparum malaria sensitive to chloroquine but resistant to pyrimethamine/sulfadoxine in Sennar, Sudan*. Trans R Soc Trop Med Hyg, 1991. 85(4): p. 446.
52. Bayoumi, R.A., H.A. Babiker, S.M. Ibrahim, H.W. Ghalib, B.O. Saeed, S. Khider, M. Elwasila, and E.A. Karim, *Chloroquine-resistant Plasmodium falciparum in eastern Sudan*. Acta Trop, 1989. 46(3): p. 157-65.