

**EDITORIAL**

**Production and Characterization of Glucosamine from Bovine Synovial Fluid**

Anwar <sup>1</sup>, M A, and El-hadi <sup>1\*</sup> M M A

<sup>1</sup>Department of Chemistry and Pharmacognosy, Faculty of Pharmacy, University of

Gezira, Wad Medani, Sudan

*Corresponding author:* Anwar, MA

*This article has been published in the GJHS Dec 2008; Vol 4 (2): 80-87. It is presented again because of previous loss of some information due to printing mistake.*

**Abstract:**

**Objectives:** Glucosamine is a natural aminomonosaccharide, which is a normal constituent of glycosaminoglycans in cartilage matrix and synovial fluid. Glucosamine has a role in the synthesis of cell membrane, collagen, osteoid, and bone matrix.

The present work was undertaken to investigate bovine synovial fluid for the production of glucosamine. The method used for isolation of glucosamine included acid hydrolysis of the fluid.

**Results:** The produced glucosamine (0.73-2.60%) had been purified and characterized by different identification methods that include: chemical colour tests; thin layer chromatography; C<sub>18</sub>- high performance liquid chromatography and infrared spectrophotometry compared to standard glucosamine.

We can conclude that, there is an essential need for every body to produce and/or to take glucosamine supplements to stimulate the production of synovial fluid which lubricates cartilage and keeps joint healthy. This led to the fact that reliable analytical methods should be adopted for quality assessment of glucosamine commercial products used by consumers. In this research the investigated bovine synovial fluid was recognized to afford a considerable amount of glucosamine obtained by acid hydrolysis.

**Conclusions:** The adopted and the described methods of analysis could be applied in qualitative and quantitative determination surveys of different supplement products containing glucosamine.

**الخلاصة**

يعتبر الجلوكوزامين (Glucosamine) من السكريات الطبيعية احادية الامين وهو المكون الاساسي لماده الجلايكوزامينوجلايكان (Glycosaminoglycans) الموجوده في الوسائد الغضروفية والوسائل السينوفي . ويلعب الجلوكوزامين دورا في تصنيع الغشاء المبطن للخلايا والكولاجين والعظام.

أجريت هذه الدراسة لإنتاج الجلوكوزامين من السائل السينوفي البقري . تم فصل الجلوكوزامين (0.73-2.60%) عن طريق التحليل المائي بواسطة حمض الكبريتيك وحمض الهيدروكلوريك المركزين. كما تم التعرف علي الجلوكوزامين المنتج بواسطة

## **EDITORIAL**

الاختبارات الكيميائية اللونية, كروماتوغرافيا الطبقة الرقيقة, وكروماتوغرافيا السوائل عاليه الكفاءه ومطيافيه الاشعه تحت الحمراء وذلك بمقارنته مع ماده المرجعيه.

يمكن أن نستنتج ان للجلوكوزامين حوجه ماسه لأجسامنا ويمكن تناوله كمكمل غذائي لتحفيز تصنيع السائل السينوفي الذي يلين الغضروف ويكسب المفاصل صحة جيدة، لذلك وفي هذا البحث وبعد الاستقصاء وجد ان السائل السينوفي البقري والمعالج بالتحليل المائي الحمضي ينتج كميته معتبره من الجلوكوزامين. كما ان الطرق المستخدمه والموصوفه للتحليل الكيميائي يمكن تطبيقها كميًا ونوعيًا لتحديد نسبه الجلوكوزامين في مختلف المنتجات المحتويه عليه.

**Key words:** Glucosamine production; Characterization; Glycosaminoglycans; Glucoproteins Bovine Synovial Fluid.

## **Introduction:**

Glucosamine occurs naturally in bacteria, plants, and animals synovial fluid; vitreous humor of the eyes; articular cartilage; umbilical cord; cornea and skin. The exoskeleton of crustaceans, cuticle of insects and cell walls of fungi also contain glucosamine as their main component is chitin (Nguyen, 2002).

Glucosamine in chitins-the main commercial source-is either pre-existent as glucoproteins or as glucosamine glucoside and/or formed in the process of hydrolysis by the interaction of carbohydrate residues and amino acids (Winifred, 1915).

Bovine synovial fluid, the whole weight of complex is accounted for in terms of protein, acetyl glucosamine, glucuronic acid and mineral content (Ogston and Stanier, 1952).

The major methods used for glucosamine production are: Acid hydrolysis and enzymes hydrolysis, although, precipitation by centrifugation and ultrafiltration may sometimes precede acid or enzyme hydrolysis (Ogston and Stanier, 1950; Ogston and Stanier, 1952).

## **Materials and Methods:**

### **Materials**

#### **Synovial Fluid**

Bovine synovial fluid was obtained from a local slaughter house. The fluid was carefully withdrawn from the knee joint with a syringe through puncture needle, 15-30minutes after slaughtering.

### **Methods**

#### **Hydrolysis of Synovial Fluid with Concentrated Sulfuric Acid and/or Hydrochloric Acid**

As described by Winifred (1915), 150 grams of the synovial fluid were treated with 100ml of concentrated sulphuric acid (a) and/or hydrochloric acid (b), and refluxed on water bath for four hours. The solution was then allowed to cool for another four hours, diluted with 200ml water and allowed to stand for 24 hours.

## **EDITORIAL**

Sulphuric acid was removed by barium carbonate, while excess hydrochloric acid was removed by vacuum evaporation. The precipitated barium sulphate was filtered. The brown filtrates were decolourized by heating with charcoal and then filtered. The resulting solutions (a and b) were concentrated at room temperature and partitioned in butanol separately. Butanol fraction which proved to contain no glucosamine was rejected, while small amount of ammonia solution (just to make alkaline) was added to the water fraction of both (a) and (b) solution mixture separately. After hydrolysis, decolourization, concentration and partitioning, the obtained water fraction (a) and (b) solution was examined for glucosamine using chitosan test, Fehling's and Benedict's solution. Furthermore, fraction (a) obtained by sulphuric acid hydrolysis which afforded sufficient product was subjected to column fractionation, thin layer chromatography (TLC) tracing, C<sub>18</sub>-HPLC and FTIR analytical methods for qualitative and quantitative determination of glucosamine.

### **Confirmatory Colour Tests**

The produced hydrolyzed solution mixtures (a) and (b) were examined separately for glucosamine qualitative chemical colour tests with chitosan (potassium hydroxide, 90% alcohol, iodine potassium iodide and dilute sulphuric acid); Fehling's (CuSO<sub>4</sub>. 5H<sub>2</sub>O + Sodium potassium tartarate); Benedict's (Na citrate, Na carbonate and CuSO<sub>4</sub>) and ninhydrin.

### **Column Chromatography**

Glass columns of 2.0 cm diameter were used. Columns were prepared using 12 grams silica gel of mesh size (0.063 – 0.2 mm) for column chromatography by using dry packing method. The water fraction of the hydrolyzed product (a) was dissolved in water (1-2ml) and added to the top of a dry-packed column. Flash column chromatography elution commenced with 50ml of water and water/ethanol (10 ml) mixture in a decreasing manner as shown in Table 1. Fractions of 10 ml were collected and analyzed by TIC using butanol, acetic acid and water (BAW) in a ratio of (3:1:1) as solvent system, ninhydrin and iodine vapour were used as locating reagents.

**Table 1: Column chromatography of the water fraction of the hydrolyzed Bovine synovial fluid.**

<b>Fraction number</b>	<b>Eluent</b>	<b>Amount of eluent used</b>
1 – 5	Water	50 ml
6	Water/ethanol (7:3)	10 ml
7	Water/ethanol (5:5)	10 ml
8	Water/ethanol (3:7)	10 ml
9	Ethanol	10 ml

## **EDITORIAL**

Similar fractions were mixed together. TLC results showed four distinct pink spots one of which (fraction 5 and 6) was corresponding to glucosamine standard sample ( $R_f = 0.51$ ).

### **Layer Chromatography (TLC)**

TLC plates were prepared using silica gel G type 60 with  $\text{CaSO}_4$  as binder. A slurry of silica gel in water was applied by means of a spreader to glass plates, previously cleaned with ethanol. A layer of 0.25 mm thick was used routinely and the plates were activated by heating at 100-110°C in an oven for an hour.

The produced hydrolyzed solution mixtures (a and b) together with the standard glucosamine were co-chromatographed on TLC plates, using butanol, acetic acid and water (BAW) in a ratio of (3:1:1) as solvent system, ninhydrin and iodine vapour were used as locating reagent.

### **C<sub>18</sub>-High Performance Liquid Chromatography (C<sub>18</sub> - HPLC)**

The produced hydrolyzed solution mixture (a) and glucosamine standard sample were co-chromatographed on C<sub>18</sub>-HPLC column ODS 4.6 mm× 150mm 5 mic (HPLC Sykam; S3210UV/visible detector; Germany). Acetonitrile and water (10:90) were used as mobile phase at a flow rate of 1ml/min at ambient temperature. UV detector was operated at 240nm.

In this method 500mg of the dried sample product (fraction 5 and 6) were dissolved in 250ml of the mobile phase, filtered and sonicated for 10 minutes. 250mg of the standard glucosamine sulphate was dissolved in 100ml mobile phase, filtered and sonicated for 10 minutes. 20µl from each was injected into the HPLC system and the peaks monitored by UV absorbance at 240nm.

### **Infrared Spectroscopy**

IR spectra of the produced compound (fraction 5 and 6) and reference glucosamine sulphate were recorded on FTIR spectrophotometer (Thermo Nicolet FTIR spectrophotometry, USA), using KBr disk of IR spectroscopic grade (Central Drug House Laboratory Reagents).

### **Optical Rotation**

The optical activity of glucosamine sulphate isolated was measured on a polarimeter (Bellingham, Stanley Limited, England).


### **Results and Discussion:**

Glucosamine and its derivative N-acetylglucosamine are synthesized in all organisms, including bacteria, yeast, fungi, plants and animals. In humans, glucosamine and N-acetylglucosamine are precursors of the disaccharide units in glycosaminoglycans which are necessary to repair and maintain healthy cartilage and joint function (Deng, 2004).



## **EDITORIAL**

Chitin hydrolysis is considered as the major natural commercial source of glucosamine (Deng, 2004; Zhu *et al.*, 2005). However, on hydrolysis, bovine synovial fluid and hyaluronic acid from different origin give remarkable amount of the aminosugar glucosamine (Ogston and Stanier, 1952; Curtain, 1955; Johnston, 1955; Preston *et al.*, 1965).

Among hydrolysis procedures, acid hydrolysis of synovial fluid and/or hyaluronic acid is the most common preferable mechanism for production of glucosamine (Zhu *et al.*, 2005). However, glucosamine hydrochloride was readily formed in a total yield of 3.4% when *Boletus edulis* fungi had been hydrolyzed by Winifred (1915) with concentrated hydrochloric acid and who concluded that glucosamine is derived from a complex nature of glucoproteins, carbohydrate and amino acid residues. Likewise, for synovial fluid being a glucoprotein its acid hydrolysis can follow the same pattern of reactions as postulated by Winifred (1915) can be summerized as follows:



**OR**




Therefore, it can be concluded that degradation of hyaluronic acid could be achieved under acid conditions to give a mixture of products one of which is the aminosugar, glucosamine which is known to be resistant to further acid hydrolysis (Kaye and Stacey, 1950).


In this study, chemical tests for the presence of glucosamine in the produced hydrolyzed mixtures (a) and (b) was confirmed when its characteristic identifying colour tests were carried out where both mixtures readily reduced Fehling's solution, gave positive chitosan test, produced yellowish brown precipitate with Benedict's and gave four distinct pink coloured spots with ninhydrin on comparative TLC chromatogram where the dominant spot of which corresponds to glucosamine standard with R<sub>f</sub> value = 0.51.

When the obtained solution mixtures of (a) and (b) are partitioned separately in butanol, glucosamine appeared as dominating compound in water fraction while butanol layer was found to be devoid of or containing very low amount as confirmed by comparative TLC.

**EDITORIAL**



EDITORIAL



- a. Standard, glucosamine sulphate.
- b. Sample (Test), glucosamine sulphate.

**Figure 1: C<sub>18</sub> – HPLC chromatogram.**

In flash column chromatography, the water fraction which is rich in glucosamine and obtained by sulphuric acid hydrolysis (a), was further fractionated on silica gel for column chromatography. Separation was initiated with water; water/ethanol mixed solvent with gradual increase in ethanol content was used to perform further separation. Glucosamine-containing fractions 5 and 6 were eluted from the mixture with water and water/ethanol (7:3), yielding 0.73 – 2.60% yellowish - white product which readily reduced Fehling's solution, gave positive chitosan test, produced yellowish brown precipitate with Benedict's and gave a pink coloured single spot with ninhydrin on comparative TLC chromatogram ( $R_f = 0.51$ ). Permanent rotation of the compound in water,  $[\alpha]_D = +70.0 - +73.5^\circ$ , which corresponds to that determined by Winifred (1915).

In reversed – phase C<sub>18</sub>-HPLC comparative analysis performed at UV 240nm and acetonitrile and water (10:90) were used as mobile phase, a peak of the same retention time (2.483minutes), appeared in chromatograms of both tested solution and the standard (Fig. 1). The substance was undoubtedly glucosamine sulphate, which was further confirmed by FTIR analysis.


The yield of glucosamine obtained from bovine synovial fluid (0.73 – 2.60%) seems to be remarkably higher compared with the amount (0.012 – 0.028%) obtained by Ogston and Stanier, (1952) and the amount (0.037 – 0.043%) obtained by Curtain, (1955), while chitin of *B.edulis* yield was 3.4% glucosamine, as reported by Winifred, (1915) from a similar process of hydrolysis.

Infrared spectra of glucosamine sulphate sample and standard were obtained using FTIR instrument (Thermo Nicolet FTIR spectrophotometer, USA) for the frequency range 4000 – 400  $cm^{-1}$ , (Fig. 2 and Fig. 3).


**EDITORIAL**

It was noted that compounds containing C-H, N-H and O-H group have absorption bands in the frequency ranges as: C – H, 3100 – 2850  $cm^{-1}$ ; N–H, 3500 – 3150  $cm^{-1}$ ; O – H, 3600 – 3200  $cm^{-1}$  (Gunstone and Sharp, 1970; Shriner *et al.*, 1980; Dudley and Felming, 1980).

Fig.2 and Fig.3 show that, characteristic and diagnostic bands of glucosamine are visible at 3347  $cm^{-1}$ , 1382  $cm^{-1}$ , 1126  $cm^{-1}$  and 615  $cm^{-1}$ . Literature agrees in assigning 3347  $cm^{-1}$  band to –NH<sub>2</sub>, -OH and C-H stretching vibration. In addition lower IR finger print region of the isolated glucosamine was found to be identical to that of the reference sample. Moreover, as the –NH<sub>2</sub>, -OH and C-H stretching



**Figure 3: FTIR Spectra of solid glucosamine sulphate (Test)**



## **EDITORIAL**

### **Figure 4 : FTIR Spectra of glucosamine sulphate , standard ( \_\_\_\_\_ ) and glucosamine sulphate sample (Test):** ° ° °

bands are aligned, thus appear as a characteristic broad band from (3600-2750  $cm^{-1}$ ) in the spectrum. The band at 1126  $cm^{-1}$  represents the free amino group (-NH<sub>2</sub>) at C<sub>2</sub> position of glucosamine. The peak at 1382  $cm^{-1}$  represents the -C-O stretching of primary alcohol group (-CH<sub>2</sub>OH) and the peak at 615  $cm^{-1}$  representing -N-H bending band (Gunstone *et al.*, 1970; Shriner *et al.*,1980; Dudley and Felming,1980; Saraswathy *et al.*,2001;Guo *et al.*,2002;Tanveer *et al.*,2002; Davika and Varsha.,2006).

In short, infrared spectra for both the isolated and the standard sample reproduce those shown in literature confirming that the isolated compound was indeed glucosamine sulphate.

### **Conclusions:**

Dietary supplements containing glucosamine are popular on the local market and many of them may not provide high quality material or need labeled quantities . This led to the fact that reliable analytical methods should be carried for quality assessment of glucosamine commercial products used by consumers. In this research the investigated bovine synovial fluid was recognized to afford a considerable amount of glucosamine obtained by acid hydrolysis. Moreover the adopted and previously described methods of analysis could be applied in qualitative and quantitative determination surveys of different supplement products containing glucosamine.

### **Acknowledgement**

We wish to thank Dr. Ayman, YW. Department of Pharmaceutics, Faculty of Pharmacy, University of Gezira for HPLC analysis and Dr. Mahmoud, E. Custom and Duties Laboratory, Khartoum for IR analysis.

### **References:**

1. Curtain, CC (1955). The nature of the protein in the Hyaluronic complex of Bovine Synovial fluid. *Biochem. J.* 61 (4): 688 – 697.
2. Davika, RB and Varsha, BP (2006). Studies on effect of pH on Cross-linking of chitosan with sodium tripoly phosphate: A Technical Note. *Pharm Sci Tech.* 7 (2): Article 50.
3. Deng, MD (2004). Metabolic engineering for production of glucosamine and N-acetyl glucosamine. Industrial Microbiology Annual Meeting; San Diego, California USA.
4. Dudley, HW and Felming (1980). Textbook. Spectroscopic methods in organic chemistry. Third edition,

**EDITORIAL**

- McGraw-Hill book company U.K.
5. Gunstone, FD and Sharp, JT (1970). An introductory course in practical organic chemistry. First edition, William Clowes and Co Ltd, London, U.K.
  6. Guo, ZC; Han, L; Hu, B and Li, SN (2002). Spectra characteristics of the complexes of glucosamine and carboxymethyl glucosamine with ferrum (II), Zinc (II), Cobalt (II) and Copper (II). *Guang Pu Xue Yu Ginag Pu FenXi*, 22 (6): 963 – 6.
  7. Johnston, JP (1955). A comparison of the properties of hyaluronic acid from normal and pathological human synovial fluids. *Biochem. J.* 59: 626 – 633.
  8. Kaye, AG and Stacey, M (1950). Observations on the chemistry of hyaluronic acid. The 283rd Meeting of the Biochemical Society, London, U.K.
  9. Nguyen, VH (2002). Valorization of chitin with the aim of production of biofungicides. VAR. Veterinary and Agrochemical Research center.
  10. Ogston, AG and Stanier, JE (1950). On the state of hyaluronic acid in synovial fluid. *Biochem. J.* 46: 364 – 376.
  11. Ogston, AG and Stanier, JE (1952). Further observations on the preparation and composition of the hyaluronic acid complex of ox synovial fluid. *Biochem. J.* 52(1): 149 – 156.
  12. Preston, BN; Davies, M and Ogston, AG (1965). The composition and physicochemical properties of hyaluronic acid prepared from ox synovial fluid and from a case of Mesothelioma. *Biochem. J.* 96: 449 – 471.
  13. Saraswathy, G; Pal, S; Rose, C and Sastry, TP (2001). A novel bio-inorganic bone implant containing deglued bone, chitosan and gelatin. *Bull Mater. Sci*, 24: 415 – 420.
  14. Shriner, RL; Fuson, RC; Curtin, DY and Morrill, TC (1980). The systematic identification of organic compounds. Six edition, John Wiley and Sons, New York, USA.
  15. Tanveer, AK; Kok, KP and Ch'ng, SH (2002). Reporting degree of deacetylation values of chitosan: The influence of analytical methods. *J Pharmaceutical Sci*, 5 (3): 205 – 212.
  16. Winifred, R (1915). The origin of the glucosamine obtained in the hydrolysis of *Boletus edulis*. Chemical Research laboratory, United College of St Salvator and St Leonard, University of St Andrews. *Biochem. J.* 313-319.
  17. Zhu, X; Cai, J; Yang, J and Su, Q (2005). Determination of glucosamine in impure chitin samples by high performance liquid chromatography. *Carbohydrate research*. 340: 1732 – 1738.